

Functions, relations and graphs


In graph sketching, one must find x and y -intercepts, show asymptotic behaviour and, determine location of stationary points.

Graphs of functions defined by

$$f(x) = ax^m + \frac{b}{x^n} \text{ for } m, n = 1, 2$$

Use the method of addition of ordinates to sketch this type of functions.

Example 1 Sketch $y = \frac{1}{2}x + \frac{1}{x}$.



Clearly the function has no axis intercepts. It shows asymptotic behaviour:

As $x \rightarrow 0^-$, $y \rightarrow -\infty$. As $x \rightarrow 0^+$, $y \rightarrow +\infty$.

$\therefore x = 0$ is an asymptote (vertical) of the function.

As $x \rightarrow -\infty$, $y \rightarrow \frac{1}{2}x$ (from below).

As $x \rightarrow +\infty$, $y \rightarrow \frac{1}{2}x$ (from above).

$\therefore y = \frac{1}{2}x$ is an asymptote (oblique) of the function.

Use calculus to find the coordinates of the stationary points:


$$y = \frac{1}{2}x + \frac{1}{x}, \frac{dy}{dx} = \frac{1}{2} - \frac{1}{x^2} = 0, \therefore x = \pm\sqrt{2}.$$

$$\text{At } x = -\sqrt{2}, y = \frac{1}{2}(-\sqrt{2}) + \frac{1}{-\sqrt{2}} = -\sqrt{2}. \text{ At } x = \sqrt{2},$$

$$y = \frac{1}{2}(\sqrt{2}) + \frac{1}{\sqrt{2}} = \sqrt{2}.$$

The stationary points are $(-\sqrt{2}, -\sqrt{2}), (\sqrt{2}, \sqrt{2})$.

Example 2 Sketch $y = x - \frac{8}{x^2}$.



x -intercept: Let $y = 0$, $x - \frac{8}{x^2} = 0$, $x^3 = 8$, $\therefore x = 2$.

Asymptotic behaviour:

As $x \rightarrow 0^-$, $y \rightarrow -\infty$. As $x \rightarrow 0^+$, $y \rightarrow -\infty$.

$\therefore x = 0$ is an asymptote of the function.

As $x \rightarrow -\infty$, $y \rightarrow x$ (from below).

As $x \rightarrow +\infty$, $y \rightarrow x$ (from below).

$\therefore y = x$ is an asymptote of the function.

Stationary points:


$$y = x - \frac{8}{x^2}, \frac{dy}{dx} = 1 + \frac{16}{x^3} = 0, \therefore x = \sqrt[3]{-2^4} = -2\sqrt[3]{2}.$$

$$\text{At } x = -2\sqrt[3]{2} = -2.52,$$

$$y = -2\sqrt[3]{2} - \frac{8}{\left(-2\sqrt[3]{2}\right)^2} = -2\sqrt[3]{2} - 2\sqrt[3]{2} = -3.78.$$

The stationary point is $(-2.52, -3.78)$.

Example 3 Sketch $y = \frac{1}{2}x^2 + \frac{2}{x}$.



x -intercept: Let $y = 0$, $\frac{1}{2}x^2 + \frac{2}{x} = 0$, $x^3 = -4$, $\therefore x = -1.59$.

Asymptotic behaviour:

As $x \rightarrow 0^-$, $y \rightarrow -\infty$. As $x \rightarrow 0^+$, $y \rightarrow +\infty$.

$\therefore x = 0$ is an asymptote of the function.

As $x \rightarrow -\infty$, $y \rightarrow \frac{1}{2}x^2$ (from below).

As $x \rightarrow +\infty$, $y \rightarrow \frac{1}{2}x^2$ (from above).

Note: $y = \frac{1}{2}x^2$ is not labeled as an asymptote. Asymptotes are straight lines.


Stationary points:

$y = \frac{1}{2}x^2 + \frac{2}{x}$, $\frac{dy}{dx} = x - \frac{2}{x^2} = 0$, $\therefore x = \sqrt[3]{2} = 2^{\frac{1}{3}}$.

At $x = 2^{\frac{1}{3}} = 1.26$, $y = \frac{1}{2}\left(2^{\frac{1}{3}}\right)^2 + \frac{2}{2^{\frac{1}{3}}} = 2^{\frac{1}{3}} + 2^{\frac{2}{3}} = 2.38$.

The stationary point is $(1.26, 2.38)$.

Example 4 Sketch $y = -\frac{x^2}{4} + \frac{4}{x^2}$.



x -intercept: Let $y = 0$, $-\frac{x^2}{4} + \frac{4}{x^2} = 0$, $x^4 = 16$, $\therefore x = \pm 2$.

Asymptotic behaviour:

As $x \rightarrow 0^-$, $y \rightarrow +\infty$. As $x \rightarrow 0^+$, $y \rightarrow +\infty$.

$\therefore x = 0$ is an asymptote of the function.

As $x \rightarrow -\infty$, $y \rightarrow -\frac{x^2}{4}$ (from above).

As $x \rightarrow +\infty$, $y \rightarrow -\frac{x^2}{4}$ (from above).

The function has no stationary points.

Graphs of functions defined by $f(x) = \frac{1}{ax^2 + bx + c}$

The sketching method of $f(x) = \frac{1}{ax^2 + bx + c}$ depends on the linear factors of $ax^2 + bx + c$.

Case 1 $ax^2 + bx + c$ can be factorised, i.e. the discriminant $b^2 - 4ac > 0$.

$$\therefore f(x) = \frac{1}{a(x-p)(x-q)}$$

There are two vertical asymptotes, $x = p$ and $x = q$.

Example 1

Sketch the graph of the function $f(x) = \frac{1}{2x^2 - x - 1}$.

Check: $b^2 - 4ac > 0$. Factorise the denominator:

$$f(x) = \frac{1}{2x^2 - x - 1} = \frac{1}{2\left(x + \frac{1}{2}\right)(x-1)}$$

Asymptotic behaviour:

As $x \rightarrow -\frac{1}{2}$ (from the left), $y \rightarrow +\infty$.

As $x \rightarrow -\frac{1}{2}$ (from the right), $y \rightarrow -\infty$.

$\therefore x = -\frac{1}{2}$ is an asymptote (vertical).

As $x \rightarrow 1$ (from the left), $y \rightarrow -\infty$.

As $x \rightarrow 1$ (from the right), $y \rightarrow +\infty$.

$\therefore x = 1$ is an asymptote (vertical).

As $x \rightarrow -\infty$, $y \rightarrow 0$ (from above).

As $x \rightarrow +\infty$, $y \rightarrow 0$ (from above).

$\therefore y = 0$ is an asymptote (horizontal).


y -intercept: Let $x = 0$, $y = f(0) = -1$.

The function has no x -intercepts.


Stationary points: $y = \frac{1}{2x^2 - x - 1}$,

$$\frac{dy}{dx} = -\frac{4x-1}{(2x^2-x-1)^2} = 0, \therefore x = \frac{1}{4}; y = -\frac{8}{9}$$

Stationary point is $\left(\frac{1}{4}, -\frac{8}{9}\right)$.



Alternatively, sketch the graph of $y = 2x^2 - x - 1$ first and then the reciprocal $y = \frac{1}{2x^2 - x - 1}$.



Case 2 $ax^2 + bx + c$ has the discriminant $b^2 - 4ac = 0$. It has two linear factors and they are the same.

$\therefore f(x) = \frac{1}{a(x-d)^2}$, and it is the transformation (dilation and translation) of $f(x) = \frac{1}{x^2}$.

Example 1 Sketch $y = \frac{1}{2x^2 - 4x + 2}$.

Check: $b^2 - 4ac = 0$. Factorise the denominator:

$$y = \frac{1}{2(x-1)^2}$$

As $x \rightarrow 1$ (from the left), $y \rightarrow +\infty$.

As $x \rightarrow 1$ (from the right), $y \rightarrow +\infty$.

$\therefore x = 1$ is an asymptote (vertical).


As $x \rightarrow -\infty$, $y \rightarrow 0$ (from above).

As $x \rightarrow +\infty$, $y \rightarrow 0$ (from above).


$\therefore y = 0$ is an asymptote (horizontal).

y-intercept: Let $x = 0$, $y = \frac{1}{2}$.

The function has no x-intercepts and stationary points.



Alternatively, sketch the graph of $y = 2x^2 - 4x + 2$ first and then the reciprocal $y = \frac{1}{2x^2 - 4x + 2}$.




Case 3 $ax^2 + bx + c$ cannot be factorised, i.e. the discriminant $b^2 - 4ac < 0$.

In this case, the quadratic is never zero and therefore, $f(x)$ has no vertical asymptotes.

The best way to sketch $f(x)$ is to sketch the quadratic first and then its reciprocal.

Example 1 Sketch $f(x) = \frac{1}{x^2 + 2x + 3}$.


Check: $b^2 - 4ac < 0$. No linear factors.



Graphs of ellipses

The general equation of an ellipse in Cartesian form with the centre at the origin is $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, where $\pm a$ are the x-intercepts and $\pm b$ are the y-intercepts.

If the ellipse is translated so that its centre is at (h, k) , the general equation becomes $\frac{(x-h)^2}{a^2} + \frac{(y-k)^2}{b^2} = 1$, where $[-a+h, a+h]$ is the domain and $[-b+k, b+k]$ the range of the relation.



Graphs of hyperbolas

The general equation of a hyperbola with its centre at the origin is $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$ or $\frac{x^2}{a^2} - \frac{y^2}{b^2} = -1$, where $\pm a$ are the x -intercepts for the first equation and $\pm b$ are the y -intercepts for the second. Both relations have two oblique asymptotes given by $y = \pm \frac{b}{a}x$.

The domain for the first is $R \setminus (-a, a)$ and the range is R . The domain for the second is R and the range is $R \setminus (-b, b)$.

If the hyperbola is translated so that its centre is at (h, k) , the general equation becomes

$$\frac{(x-h)^2}{a^2} - \frac{(y-k)^2}{b^2} = 1 \quad \text{or} \quad \frac{(x-h)^2}{a^2} - \frac{(y-k)^2}{b^2} = -1.$$

The two oblique asymptotes for both relations are

$$y - k = \pm \frac{b}{a}(x - h).$$


The domain for the first is $R \setminus (-a+h, a+h)$ and the range is R . The domain for the second is R and the range is $R \setminus (-b+k, b+k)$.

Example 1 Sketch the graphs of

a) $x^2 + \frac{y^2}{9} = 1$ b) $(x+1)^2 + \frac{(y-2)^2}{9} = 1$.

Show the x , y intercepts, the domains and ranges.

a) x -intercepts: $x = \pm 1$; y -intercepts: $y = \pm 3$




Domain: $[-1, 1]$; range: $[-3, 3]$.

b) x -intercepts: Let $y = 0$, $(x+1)^2 + \frac{4}{9} = 1$, $(x+1)^2 = \frac{5}{9}$,



$$x+1 = \pm \frac{\sqrt{5}}{3}, \quad x = -1 \pm \frac{\sqrt{5}}{3}.$$

y -intercepts: Let $x = 0$, $1 + \frac{(y-2)^2}{9} = 1$, $y = 2$.

Centre: $(-1, 2)$.



Domain: $[-2, 0]$; range: $[-1, 5]$.




Example 1 Sketch the graphs of a) $\frac{x^2}{9} - \frac{y^2}{4} = 1$ and

b) $\frac{(x+2)^2}{9} - \frac{y^2}{4} = -1$.

State the x and y intercepts, domains and ranges, and equations of the asymptotes.

a) x -intercepts: $x = \pm 3$; no y -intercepts.


Equations of asymptotes: $y = \pm \frac{2}{3}x$.



Domain: $R \setminus (-3, 3)$; range: R .

b) y -intercepts: Let $x = 0$, $\frac{4}{9} - \frac{y^2}{4} = -1$, $y = \pm \frac{2\sqrt{13}}{3}$; no x -intercepts.

Equations of asymptotes: $y = \pm \frac{2}{3}(x+2)$.



Domain: R ; range: $R \setminus (-2, 2)$.

Reciprocal trigonometric functions

The cosecant function of x , $\operatorname{cosec} x$, is defined as the reciprocal of the sine function, $\sin x$, i.e.


$$\operatorname{cosec} x = \frac{1}{\sin x}.$$

The other reciprocal functions are defined as

$$\sec x = \frac{1}{\cos x} \quad \text{and} \quad \cot x = \frac{1}{\tan x}.$$

Graphs of $\operatorname{cosec} x$, $\sec x$ and $\cot x$

$y = \operatorname{cosec}(x)$




Asymptotes: $x = n\pi$, where $n = 0, \pm 1, \pm 2, \dots$

Domain: $R \setminus \{x : x = n\pi, n = 0, \pm 1, \pm 2, \dots\}$;

range: $(-\infty, -1] \cup [1, \infty)$.

Period: 2π .

$y = \sec(x)$




Asymptotes: $x = n\pi$, where $n = \pm \frac{1}{2}, \pm \frac{3}{2}, \pm \frac{5}{2}, \dots$

Domain: $R \setminus \left\{x : x = n\pi, n = \pm \frac{1}{2}, \pm \frac{3}{2}, \pm \frac{5}{2}, \dots\right\}$;

range: $(-\infty, -1] \cup [1, \infty)$.

Period: 2π .

$y = \cot(x)$



Asymptotes: $x = n\pi$, where $n = 0, \pm 1, \pm 2, \dots$

Domain: $R \setminus \{x : x = n\pi, n = 0, \pm 1, \pm 2, \dots\}$; range: R .


Period: π .

Transformations of cosec x , sec x and cot x

The transformations-dilation, reflection and translation are applicable to all functions. Always carry out translation last.

Example 1 Sketch $y = -\sec(2x) + 1$.

The graph of $y = \sec(x)$ is reflected in the x -axis, dilated horizontally by a factor of $\frac{1}{2}$, and translated upwards by a unit.



Asymptotes: $x = n\pi$, where $n = \pm\frac{1}{4}, \pm\frac{3}{4}, \pm\frac{5}{4}, \dots$

Domain: $R \setminus \left\{ x : x = n\pi, n = \pm\frac{1}{4}, \pm\frac{3}{4}, \pm\frac{5}{4}, \dots \right\}$;

range: $(-\infty, 0] \cup [2, \infty)$.

Period: π .


Example 2 Sketch $y = \cot\left(\frac{\pi}{2} - x\right)$ and $-\pi \leq x \leq \pi$.

Rewrite: $y = \cot\left(\frac{\pi}{2} - x\right) = \cot\left[-\left(x - \frac{\pi}{2}\right)\right]$.

The graph of $y = \cot(x)$ is reflected in the y -axis, and then

translated to the right by $\frac{\pi}{2}$.

The graph is restricted to $-\pi \leq x \leq \pi$.



Asymptotes: $x = -\frac{\pi}{2}$ and $x = \frac{\pi}{2}$.

Domain: $[-\pi, \pi] \setminus \left\{ -\frac{\pi}{2}, \frac{\pi}{2} \right\}$; range: R .

Period: π .

Identities

An identity is an equality which is always true as long as it is defined.

$$\tan x = \frac{\sin x}{\cos x}$$

$$\cot x = \frac{\cos x}{\sin x}$$

$$\sin^2 x + \cos^2 x = 1$$

$$\cos^2 x = 1 - \sin^2 x$$

$$\sin^2 x = 1 - \cos^2 x$$

$$\sec^2 x = 1 + \tan^2 x$$

$$\operatorname{cosec}^2 x = 1 + \cot^2 x$$

Example 1 Use $\sin^2 x + \cos^2 x = 1$ to prove $\sec^2 x = 1 + \tan^2 x$ and $\operatorname{cosec}^2 x = 1 + \cot^2 x$.

$$1 = \cos^2 x + \sin^2 x, \quad \frac{1}{\cos^2 x} = \frac{\cos^2 x + \sin^2 x}{\cos^2 x},$$

$$\frac{1}{\cos^2 x} = \frac{\cos^2 x}{\cos^2 x} + \frac{\sin^2 x}{\cos^2 x}, \quad \therefore \sec^2 x = 1 + \tan^2 x.$$

$$1 = \sin^2 x + \cos^2 x, \quad \frac{1}{\sin^2 x} = \frac{\sin^2 x + \cos^2 x}{\sin^2 x},$$

$$\frac{1}{\sin^2 x} = \frac{\sin^2 x}{\sin^2 x} + \frac{\cos^2 x}{\sin^2 x}, \quad \therefore \operatorname{cosec}^2 x = 1 + \cot^2 x.$$

Example 2 Prove $\frac{1}{1 - \sin x} = \sec x(\sec x + \tan x)$.

$$RHS = \frac{1}{\cos x} \left(\frac{1}{\cos x} + \frac{\sin x}{\cos x} \right) = \frac{1}{\cos x} \left(\frac{1 + \sin x}{\cos x} \right)$$

$$= \frac{1 + \sin x}{\cos^2 x} = \frac{1 + \sin x}{1 - \sin^2 x} = \frac{1 + \sin x}{(1 - \sin x)(1 + \sin x)}$$

$$= \frac{1}{1 - \sin x} = LHS.$$

Compound angle formulas

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

Double angle formulas

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A$$

$$\cos 2A = 2 \cos^2 A - 1$$

$$\cos 2A = 1 - 2 \sin^2 A$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$

Example 1 Prove $\tan x + \tan y = \frac{\sin(x+y)}{\cos x \cos y}$.

$$\begin{aligned} LHS &= \frac{\sin x}{\cos x} + \frac{\sin y}{\cos y} = \frac{\sin x \cos y + \cos x \sin y}{\cos x \cos y} \\ &= \frac{\sin(x+y)}{\cos x \cos y} = RHS \end{aligned}$$

Example 2 Use the compound angle formulas for $\sin(A \pm B)$ and $\cos(A \pm B)$ to prove that for $\tan(A \pm B)$.

$$\begin{aligned} \tan(A \pm B) &= \frac{\sin(A \pm B)}{\cos(A \pm B)} = \frac{\sin A \cos B \pm \cos A \sin B}{\cos A \cos B \mp \sin A \sin B} \\ &= \frac{\sin A \cos B \pm \cos A \sin B}{\cos A \cos B \mp \sin A \sin B} = \frac{\frac{\sin A \cos B}{\cos A \cos B} \pm \frac{\cos A \sin B}{\cos A \cos B}}{\frac{\cos A \cos B}{\cos A \cos B} \mp \frac{\sin A \sin B}{\cos A \cos B}} \\ &= \frac{\frac{\sin A}{\cos A} \pm \frac{\sin B}{\cos B}}{1 \mp \frac{\sin A \sin B}{\cos A \cos B}} = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}. \end{aligned}$$

Example 3

Prove $(\sin x + \sin y)(\sin x - \sin y) = \sin(x+y)\sin(x-y)$.

$$\begin{aligned} RHS &= (\sin x \cos y + \cos x \sin y)(\sin x \cos y - \cos x \sin y) \\ &= \sin^2 x \cos^2 y - \cos^2 x \sin^2 y \\ &= \sin^2 x(1 - \sin^2 y) - (1 - \sin^2 x)\sin^2 y \\ &= \sin^2 x - \sin^2 y \\ &= (\sin x + \sin y)(\sin x - \sin y) = LHS \end{aligned}$$

Example 4 If $\sin x = \frac{1}{3}$ and $\sec y = \frac{5}{4}$, where

$$x, y \in \left[0, \frac{\pi}{2}\right], \text{ evaluate } \sin(x-y).$$

Note: x and y are in the first quadrant, $\therefore \cos x$ and $\sin y$ have positive values.

$$\sin x = \frac{1}{3}, \therefore \cos x = \sqrt{1 - \sin^2 x} = \sqrt{1 - \left(\frac{1}{3}\right)^2} = \sqrt{\frac{8}{9}} = \frac{2\sqrt{2}}{3}.$$

$$\sec y = \frac{5}{4}, \therefore \cos y = \frac{4}{5}, \text{ and}$$

$$\sin y = \sqrt{1 - \cos^2 y} = \sqrt{1 - \left(\frac{4}{5}\right)^2} = \frac{3}{5}.$$

$$\begin{aligned} \therefore \sin(x-y) &= \sin x \cos y - \cos x \sin y = \frac{1}{3} \times \frac{4}{5} - \frac{2\sqrt{2}}{3} \times \frac{3}{5} \\ &= \frac{2(2-3\sqrt{2})}{15}. \end{aligned}$$

Example 5 Find the exact value of $\cos \frac{\pi}{12}$.

$$\begin{aligned} \cos \frac{\pi}{12} &= \cos\left(\frac{\pi}{3} - \frac{\pi}{4}\right) = \cos \frac{\pi}{3} \cos \frac{\pi}{4} + \sin \frac{\pi}{3} \sin \frac{\pi}{4} \\ &= \frac{1}{2} \times \frac{1}{\sqrt{2}} + \frac{\sqrt{3}}{2} \times \frac{1}{\sqrt{2}} = \frac{1+\sqrt{3}}{2\sqrt{2}} \\ &= \frac{\sqrt{2}(1+\sqrt{3})}{4}. \end{aligned}$$

Example 6 Find the exact values of $\sin \frac{\pi}{8}$ and $\tan \frac{\pi}{8}$.

$$\begin{aligned} \text{For } \sin \frac{\pi}{8}, \text{ consider } \cos \frac{\pi}{4} &= \cos 2\left(\frac{\pi}{8}\right), \\ \therefore \frac{1}{\sqrt{2}} &= 1 - 2\sin^2\left(\frac{\pi}{8}\right), \therefore \sin^2\left(\frac{\pi}{8}\right) = \frac{2-\sqrt{2}}{4}. \end{aligned}$$

$$\text{Hence } \sin \frac{\pi}{8} = \frac{\sqrt{2-\sqrt{2}}}{2}, \text{ since } 0 < \frac{\pi}{8} < \frac{\pi}{2}.$$

$$\text{For } \tan \frac{\pi}{8}, \text{ consider } \tan \frac{\pi}{4} = \tan 2\left(\frac{\pi}{8}\right),$$

$$\therefore 1 = \frac{2 \tan \frac{\pi}{8}}{1 - \tan^2\left(\frac{\pi}{8}\right)}, \therefore \tan^2\left(\frac{\pi}{8}\right) + 2 \tan \frac{\pi}{8} - 1 = 0.$$

Use the quadratic formula to find

$$\tan \frac{\pi}{8} = \frac{-2 + \sqrt{4+4}}{2} = \sqrt{2} - 1.$$

Example 7 Prove $\cos 3\theta = 4\cos^3 \theta - 3\cos \theta$.

$$\begin{aligned} LHS &= \cos 3\theta = \cos(2\theta + \theta) = \cos 2\theta \cos \theta - \sin 2\theta \sin \theta \\ &= (2\cos^2 \theta - 1)\cos \theta - (2\sin \theta \cos \theta)\sin \theta \\ &= (2\cos^2 \theta - 1)\cos \theta - 2\sin^2 \theta \cos \theta \\ &= (2\cos^2 \theta - 1)\cos \theta - 2(1 - \cos^2 \theta)\cos \theta \\ &= 2\cos^3 \theta - \cos \theta - 2\cos \theta + 2\cos^3 \theta \\ &= 4\cos^3 \theta - 3\cos \theta = RHS \end{aligned}$$

Example 8 Solve $2 + \cos 2\beta = 3\cos \beta$ where $\beta \in [0, 2\pi]$.

$$2 + \cos 2\beta = 3\cos \beta, \quad 2 + 2\cos^2 \beta - 1 = 3\cos \beta,$$

$$2\cos^2 \beta - 3\cos \beta + 1 = 0,$$

$$(2\cos \beta - 1)(\cos \beta - 1) = 0,$$

$$\therefore \cos \beta = \frac{1}{2}, \quad \beta = \frac{\pi}{3}, \frac{5\pi}{3},$$


$$\text{or } \cos \beta = 1, \quad \beta = 0, 2\pi.$$

$$\text{The solution set is } \left\{0, \frac{\pi}{3}, \frac{5\pi}{3}, 2\pi\right\}.$$

Restricted trigonometric functions


Sin x

The function $f: \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] \rightarrow R, f(x) = \sin x$ is represented simply as $\sin x$. Its range is $[-1, 1]$. It is a one-to-one function and therefore its inverse is also a function. The inverse function is denoted as $\sin^{-1}x$. The domain of $\sin^{-1}x$ is $[-1, 1]$ and the range is $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$.




Cos x

The function $f: [0, \pi] \rightarrow R, f(x) = \cos x$ is represented by $\cos x$. Its range is $[-1, 1]$. It is also a one-to-one function and therefore its inverse is a function. The inverse function is denoted as $\cos^{-1}x$ with domain $[-1, 1]$ and range $[0, \pi]$.



Tan x


The function $f: \left(-\frac{\pi}{2}, \frac{\pi}{2}\right) \rightarrow R, f(x) = \tan x$ is represented by $\tan x$. Its range is R . It is a one-to-one function and its inverse is also a function. The inverse function is denoted as $\tan^{-1}x$ with domain R and range $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$.



Transformations of inverse trigonometric functions

Example 1 Sketch $y = -2\sin^{-1}(x+1)$. State the domain and range.


Start with the graph of $y = \sin^{-1}x$. Reflect in the x -axis, dilate vertically by a factor of 2, then translate to the left by a unit.



Domain: $[-2, 0]$; range: $[-\pi, \pi]$.

Example 2 Sketch $y = \frac{1}{2}\cos^{-1}(2x) - \frac{\pi}{4}$. State the domain and range.

Start with the graph of $y = \cos^{-1}x$, dilate vertically by a factor of $\frac{1}{2}$ and horizontally by the same factor, then translate downwards by $\frac{\pi}{4}$.




Domain: $\left[-\frac{1}{2}, \frac{1}{2}\right]$; range: $\left[-\frac{\pi}{4}, \frac{\pi}{4}\right]$.

Example 3 Sketch $y = -\tan^{-1}(2x+1) + \pi$. State the domain and range.

Rewrite $y = -\tan^{-1}(2x+1) + \pi$ as $y = -\tan^{-1}2\left(x + \frac{1}{2}\right) + \pi$.

Start with the graph of $y = \tan^{-1}x$, reflect in the x -axis, dilate horizontally by a factor of $\frac{1}{2}$, then translate to the left by $\frac{1}{2}$ and upwards by π .




Domain: \mathbb{R} ; range: $\left(\frac{\pi}{2}, \frac{3\pi}{2}\right)$.

Example 4 Sketch $y = \sin^{-1}\left(1 - \frac{x}{2}\right) - \frac{\pi}{2}$. State the domain and range.

Rewrite $y = \sin^{-1}\left(1 - \frac{x}{2}\right) - \frac{\pi}{2}$ as $y = \sin^{-1}\left(-\frac{1}{2}(x-2)\right) - \frac{\pi}{2}$.

Start with the graph of $y = \sin^{-1}x$, reflect in the y -axis, dilate horizontally by a factor of 2, then translate to the right by 2 units, and downwards by $\frac{\pi}{2}$.



Domain: $[0, 4]$; range: $[-\pi, 0]$.