

1a. $f(x) = \sqrt{x} + \frac{x}{2}$, $f(x+1) = \sqrt{x+1} + \frac{x+1}{2}$,
 $\therefore g(x) = 2f(x+1) = 2\sqrt{x+1} + x + 1$.

1b. $g(x) = 0 \therefore 2\sqrt{x+1} + x + 1 = 0$, $2\sqrt{x+1} = -(x+1)$,
 $\therefore 4(x+1) = (x+1)^2$, $4(x+1) - (x+1)^2 = 0$,
 $(x+1)[4 - (x+1)] = 0$, $(x+1)(3-x) = 0$.
Only $x = -1$ satisfies $g(x) = 0$.

2. $y = 1 + 3\log_e\left(\frac{2x-b}{a}\right)$ and $y = -2$ when $x = b$.
 $\therefore -2 = 1 + 3\log_e\left(\frac{b}{a}\right)$, $\therefore \log_e\left(\frac{b}{a}\right) = -1$, $\therefore \log_e\left(\frac{a}{b}\right) = 1$.

Hence $\frac{a}{b} = e$, $\therefore a = be$.

3. $f(x) = \frac{\log_e(ax)}{ax}$,
 $f'(x) = \frac{(ax)\left(\frac{1}{x}\right) - (a)(\log_e(ax))}{(ax)^2} = \frac{1 - \log_e(ax)}{ax^2}$.
 $f'(a^{-1}) = \frac{1 - \log_e(1)}{a^{-1}} = \frac{1}{a^{-1}} = a$.

4a. $y = f'(x) = \frac{1}{2}x + \frac{3}{2}$ for $-1 \leq x < 1$.
 $\therefore f(x) = \frac{1}{4}x^2 + \frac{3}{2}x + c$.
Given $f(-1) = -1$, $\therefore \frac{1}{4}(-1)^2 + \frac{3}{2}(-1) + c = -1$, $\therefore c = \frac{1}{4}$.

$\therefore f(x) = \frac{1}{4}x^2 + \frac{3}{2}x + \frac{1}{4}$ for $-1 \leq x < 1$.

Hence $y = \frac{1}{4}x^2 + \frac{3}{2}x + \frac{1}{4}$ for $-1 \leq x < 1$.


4b. For $-1 \leq x < 1$, if $f(-1) = 0$, then $\frac{1}{4}(-1)^2 + \frac{3}{2}(-1) + c = 0$,

$\therefore c = \frac{5}{4}$. Hence $y = \frac{1}{4}x^2 + \frac{3}{2}x + \frac{5}{4} = \frac{1}{4}(x+1)(x+5)$.

Similarly, for $x \in (-\infty, -1)$,

$y = -\frac{1}{4}x^2 + \frac{1}{2}x + \frac{3}{4} = -\frac{1}{4}(x+1)(x-3)$.

For $x \in [1, \infty)$, $y = 2x + 1$.



5a. $y = 1 + \cos\frac{x}{2}$, $x \in [0, 2\pi]$.


Coordinates of the inflection point are $(\pi, 1)$.

$\frac{dy}{dx} = -\frac{1}{2}\sin\frac{x}{2}$,

\therefore gradient of the tangent at $(\pi, 1) = -\frac{1}{2}\sin\frac{\pi}{2} = -\frac{1}{2}$.

Equation of the tangent: $y - 1 = -\frac{1}{2}(x - \pi)$, $y = -\frac{1}{2}x + \left(1 + \frac{\pi}{2}\right)$

5b.



Shaded area

$$\begin{aligned} &= \int_0^{\pi} \left\{ \left(-\frac{1}{2}x + \left(1 + \frac{\pi}{2}\right) \right) - \left(1 + \cos\frac{x}{2} \right) \right\} dx + \int_{\pi}^{2\pi} \left(1 + \cos\frac{x}{2} \right) dx - \frac{1}{2}(2)(1) \\ &= \int_0^{\pi} \left(-\frac{1}{2}x + \frac{\pi}{2} - \cos\frac{x}{2} \right) dx + \int_{\pi}^{2\pi} \left(1 + \cos\frac{x}{2} \right) dx - 1 \\ &= \left[-\frac{x^2}{4} + \frac{\pi x}{2} - 2\sin\frac{x}{2} \right]_0^{\pi} + \left[x + 2\sin\frac{x}{2} \right]_{\pi}^{2\pi} - 1 \\ &= \left(\frac{\pi^2}{4} - 2 \right) + (\pi - 2) - 1 \\ &= \frac{\pi^2}{4} + \pi - 5. \end{aligned}$$

6. $g(x) = a \sin x + b \cos x$, $g\left(\frac{\pi}{4}\right) = 2\sqrt{2}$, $g\left(-\frac{\pi}{6}\right) = -1$.

$$g\left(\frac{\pi}{4}\right) = a \sin \frac{\pi}{4} + b \cos \frac{\pi}{4} = 2\sqrt{2}, \therefore a\left(\frac{1}{\sqrt{2}}\right) + b\left(\frac{1}{\sqrt{2}}\right) = 2\sqrt{2},$$

$$\therefore a + b = 4 \dots\dots\dots(1)$$

$$g\left(-\frac{\pi}{6}\right) = a \sin\left(-\frac{\pi}{6}\right) + b \cos\left(-\frac{\pi}{6}\right) = -1,$$

$$\therefore a\left(-\frac{1}{2}\right) + b\left(\frac{\sqrt{3}}{2}\right) = -1, \therefore a - \sqrt{3}b = 2 \dots\dots\dots(2)$$

$$(1) - (2), b + \sqrt{3}b = 2, b(\sqrt{3} + 1) = 2,$$

$$\therefore b = \frac{2}{\sqrt{3} + 1} = \sqrt{3} - 1 \dots\dots\dots(3)$$

Substitute (3) in (1), $a = 4 - (\sqrt{3} - 1) = 5 - \sqrt{3}$.

7. $f'(x) = \frac{1}{1 - 6x + 9x^2} = \frac{1}{(1 - 3x)^2}$,

$$f(x) = \int \frac{1}{(1 - 3x)^2} dx = \int (1 - 3x)^{-2} dx = \frac{(1 - 3x)^{-1}}{3} + c$$

$$= \frac{1}{3(1 - 3x)} + c.$$

$$[f(x)]_{-\frac{1}{3}}^0 = \left[\frac{1}{3(1 - 3x)} + c \right]_{-\frac{1}{3}}^0 = \frac{1}{3} - \frac{1}{6} = \frac{1}{6}.$$

8a. $g : (-\infty, -3] \rightarrow R$, $g(x) = 1 - \frac{1}{2}|x + 2|$.

g is an increasing function. Its range is $(-\infty, g(-3)]$,

$$\text{i.e. } \left(-\infty, \frac{1}{2}\right].$$

Equation of $g(x)$:

$$y = 1 - \frac{1}{2}|x + 2| = 1 - \frac{1}{2}(-(x + 2)) = \frac{1}{2}x + 2$$

Equation of $g^{-1}(x)$:

$$x = \frac{1}{2}y + 2, \therefore y = 2(x - 2).$$

$$\therefore g^{-1}(x) = 2(x - 2).$$

8b. Domain of g^{-1} is the same as the range of g , i.e. $\left(-\infty, \frac{1}{2}\right]$.

9. Let $f(x) = \sqrt{\tan x}$,

$$f'(x) = \frac{\sec^2 x}{2\sqrt{\tan x}} = \frac{1}{2\sqrt{\tan x}(\cos^2 x)},$$

$$f'\left(\frac{\pi}{4}\right) = \frac{1}{2\sqrt{\tan \frac{\pi}{4}}\left(\cos \frac{\pi}{4}\right)^2} = 1.$$

$$\frac{\sqrt{\tan 1} - \sqrt{\tan \frac{\pi}{4}}}{1 - \frac{\pi}{4}} \approx f'\left(\frac{\pi}{4}\right), \therefore \sqrt{\tan 1} - \sqrt{\tan \frac{\pi}{4}} \approx 1 - \frac{\pi}{4}.$$

10a. ${}^6C_3 \left(\frac{1}{3}\right)^3 \left(\frac{2}{3}\right)^3 = 20 \left(\frac{1}{3}\right)^3 \left(\frac{2}{3}\right)^3$.

10b. Random variable X has a binomial distribution,

$$n = 6, p = \frac{1}{3}, q = \frac{2}{3}.$$


$$E(X) = np = 2, \text{Var}(X) = npq = \frac{4}{3}.$$

11a. $\frac{1}{2}p(2^{-4}) = 1, \therefore 3p = 1, p = \frac{1}{3}$.

11b. $f(0) = \frac{1}{2}p = \frac{1}{2}\left(\frac{1}{3}\right) = \frac{1}{6}$.

$$\Pr(X \leq 0) = 1 - \Pr(X > 0) = 1 - \frac{1}{2}\left(2\right)\left(\frac{1}{6}\right) = \frac{5}{6}.$$

12a.



$$\Pr(Z < 1) = 1 - \frac{1}{2}(1 - 0.68) = 0.84.$$

12b. $\mu = 72, \sigma = 6$.

$$\Pr(Z < 1) = \Pr(Z > -1), \therefore \Pr(X \geq x) = \Pr(Z < 1) = \Pr(Z > -1).$$

$$\therefore \frac{x - 72}{6} = -1, \therefore x = 66.$$

Please inform mathline@itute.com re conceptual, mathematical and/or typing errors