

Mathematics lesson – Binomial expansion algorithm (without calculator)

Free download and print from www.itute.com
Copyright 2009 © itute.com

The binomial theorem

This theorem is used to expand expression of the form $(p + q)^n$, where n is a positive integer.

$$(p + q)^n = {}^n C_0 p^n q^0 + {}^n C_1 p^{n-1} q^1 + {}^n C_2 p^{n-2} q^2 + \dots + {}^n C_n p^0 q^n$$

The coefficients ${}^n C_0, {}^n C_1, {}^n C_2, \dots$ can be determined by

$${}^n C_r = \frac{{}^n P_r}{r!}.$$

For example, ${}^7 C_3 = \frac{{}^7 P_3}{3!} = \frac{7 \times 6 \times 5}{3 \times 2 \times 1} = 35.$

Alternatively, for small n , the Pascal triangle can be used to evaluate those coefficients without using a calculator.

			1							
$n=1$			1	1						
$n=2$			1	2	1					
$n=3$			1	3	3	1				
$n=4$			1	4	6	4	1			
$n=5$			1	5	10	10	5	1		
$n=6$			1	6	15	20	15	6	1	
$n=7$			1	7	21	35	35	21	7	1

For example, in the expansion of $(p + q)^7$, ${}^7 C_0 = 1, {}^7 C_1 = 7,$
 ${}^7 C_2 = 21, {}^7 C_3 = 35, \dots$

Hence

$$\begin{aligned} (p + q)^7 &= {}^7 C_0 p^7 q^0 + {}^7 C_1 p^6 q^1 + {}^7 C_2 p^5 q^2 + {}^7 C_3 p^4 q^3 + {}^7 C_4 p^3 q^4 + {}^7 C_5 p^2 q^5 \\ &\quad + {}^7 C_6 p^1 q^6 + {}^7 C_7 p^0 q^7 \\ &= 1p^7 q^0 + 7p^6 q^1 + 21p^5 q^2 + 35p^4 q^3 + 35p^3 q^4 + 21p^2 q^5 \\ &\quad + 7p^1 q^6 + 1p^0 q^7 \end{aligned}$$

For large n , both manual methods become impractical and time consuming to evaluate the coefficients.

However, according to

$${}^n C_{r+1} = \frac{{}^n P_{r+1}}{(r+1)!} = \frac{n(n-1)(n-2)\dots(n-r+1)(n-r)}{r!(r+1)} = \frac{{}^n C_r (n-r)}{r+1},$$

successive coefficients can be generated in a shorter time by

$${}^n C_{r+1} = \frac{{}^n C_r (n-r)}{r+1}.$$

For example, from ${}^7 C_0 = 1, {}^7 C_1 = \frac{{}^7 C_0 (7-0)}{0+1} = 7,$

$${}^7 C_2 = \frac{{}^7 C_1 (7-1)}{1+1} = \frac{7 \times 6}{2} = 21, \quad {}^7 C_3 = \frac{{}^7 C_2 (7-2)}{2+1} = \frac{21 \times 5}{3} = 35,$$

$${}^7 C_4 = \frac{{}^7 C_3 (7-3)}{3+1} = \frac{35 \times 4}{4} = 35, \quad {}^7 C_5 = \frac{{}^7 C_4 (7-4)}{4+1} = \frac{35 \times 3}{5} = 21,$$

$${}^7 C_6 = \frac{{}^7 C_5 (7-5)}{5+1} = \frac{21 \times 2}{6} = 7, \quad {}^7 C_7 = \frac{{}^7 C_6 (7-6)}{6+1} = \frac{7 \times 1}{7} = 1.$$

To write down the terms in the expansion of $(p + q)^7$, follow the algorithm as shown below.

From the first term $1p^7q^0,$

the second term is $\frac{1 \times 7}{1} p^6 q^1 = 7p^6 q^1,$

the third term is $\frac{7 \times 6}{2} p^5 q^2 = 21p^5 q^2,$

the fourth term is $\frac{21 \times 5}{3} p^4 q^3 = 35p^4 q^3$ etc.

Using the same algorithm, the terms in the expansion of $(p + q)^{20}$ are:

$1p^{20}q^0,$

$$\frac{1 \times 20}{1} p^{19} q^1 = 20p^{19} q^1,$$

$$\frac{20 \times 19}{2} p^{18} q^2 = 190p^{18} q^2,$$

$$\frac{190 \times 18}{3} p^{17} q^3 = 1140p^{17} q^3,$$

$$\frac{1140 \times 17}{4} p^{16} q^4 = 4845p^{16} q^4,$$

$$\frac{4845 \times 16}{5} p^{15} q^5 = 15504p^{15} q^5 \text{ etc.}$$

Instead of evaluating, e.g. ${}^{20} C_6$ by manual arithmetic:

$${}^{20} C_6 = \frac{20 \times 19 \times 18 \times 17 \times 16 \times 15}{6 \times 5 \times 4 \times 3 \times 2 \times 1} = 38760, \text{ evaluation time will be}$$

shortened by using the algorithm: $\frac{15504 \times 15}{6} = 38760.$